



**PCT**

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
 International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>5</sup> : A01N 25/34, 47/36, 43/64 A01N 39/04		A1	(11) International Publication Number: <b>WO 93/13658</b> (43) International Publication Date: 22 July 1993 (22.07.93)
(21) International Application Number: PCT/US92/11305 (22) International Filing Date: 30 December 1992 (30.12.92) (30) Priority data: 07/817,247      6 January 1992 (06.01.92)      US 07/817,248      6 January 1992 (06.01.92)      US (60) Parent Applications or Grants (63) Related by Continuation US      07/817,247 (CIP) Filed on      6 January 1992 (06.01.92) US      07/817,248 (CIP) Filed on      6 January 1992 (06.01.92) (71) Applicant (for all designated States except US): E.I. DU PONT DE NEMOURS AND COMPANY (US/US); 1007 Market Street, Wilmington, DE 19898 (US).		(72) Inventor; and (75) Inventor/Applicant (for US only) : JACKISCH, David, Allan [US/US]; 307 Walden Road, Wilmington, DE 19803 (US). (74) Agents: COSTELLO, James, A. et al.; E.I. du Pont de Nemours and Company, Legal/Patent Records Center, 1007 Market Street, Wilmington, DE 19898 (US). (81) Designated States: AU, BB, BG, BR, CA, CS, FI, HU, JP, KP, KR, LK, MG, MN, MW, NO, NZ, PL, RO, RU, SD, UA, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, SN, TD, TG). Published With international search report.	
(54) Title: TABLET FORMULATION WITH INTERNAL DESICCANT			
(57) Abstract <p>A tablet formulation comprising a pesticide and a delivery system, the delivery system comprising a range of acid/base combinations, a dispersant, a wetting agent, polyvinylpyrrolidone, and a characterizing internal desiccant to ensure tablet stability, the tablet being selected to complement the particular class of base selected for the formulation.</p>			

BEST AVAILABLE COPY

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SK	Slovak Republic
CI	Côte d'Ivoire	LI	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	MC	Monaco	TG	Togo
DE	Germany	MG	Madagascar	UA	Ukraine
DK	Denmark	ML	Mali	US	United States of America
ES	Spain	MN	Mongolia	VN	Viet Nam
FI	Finland				

TITLE

## TABLET FORMULATION WITH INTERNAL DESICCANT

5 An effervescent tablet produces gas bubbles by the reaction between an acid and a base. Should the water level in the tablet exceed about 0.1%, the tablet will begin to lose its effervescence even though it is stored in its container. Normally, an amount of water exceeding 0.1% is required to formulate a tablet having acceptable integrity and strength.

10 One way to keep the water employed during formulation from ruining the tablet during long-term storage is to package the tablet with a desiccant. The water in the tablet diffuses out and is taken up by the desiccant. This water is held by the desiccant and so cannot cause the acid/base reaction to occur. However, the need for an external desiccant complicates design of the tablet package.

15 One such tablet formulation that does not have an internal desiccant is disclosed in WO 90/00007. Two formulations that contain an internal desiccant are disclosed in CA-A-2,013,918 and WO 88/09161. CA-A-2,013,918 discloses a tablet comprising potassium carbonate and/or potassium bicarbonate as bases and a desiccant which physically adsorbs water. WO 88/09161 discloses an effervescent tablet for cleaning dentures comprising pancreatin, an acid  
20 component, a base component, and a drying agent.

SUMMARY OF THE INVENTION

This invention comprises a tablet formulation consisting essentially of by total weight of the formulated composition:

- 25 (i) about 0.1% to 75% of a pesticide;  
(ii) about 25% to 99.9% of a delivery system characterized by a panel of components complementary to the pesticide of (i) having the following components:
- (a) about 5% to 75% of a dibasic or tribasic organic carboxylic acid  
30 or a mixture thereof;  
(b) about 5% to 75% of an ammonium, lithium, sodium or potassium carbonate or bicarbonate or a mixture thereof;  
(c) about 0.5% to 20% of a dispersant;  
(d) about 0.1% to 5% of water-insoluble cross-linked polyvinyl-  
35 polypyrrolidone;

- (e) about 0.1% to 5% of an anionic or nonionic wetting agent; and
- (f) about 1% to 20% of an internal desiccant being selected from the

group:

- (A) one or a mixture of desiccants that chemically bind water, and
  - 5 (B) one or a mixture of desiccants that physically adsorb water;
- the desiccant being (A) when (b) is potassium carbonate or potassium bicarbonate.

10 This delivery system is characterized by the inter-relationship of components (a) to (f) in the recited ranges to effect rapid disintegration of finely dispersed pesticide particles (i). Preferred ranges of the composition are 5% to 70%, more preferably 10% to 60% of the pesticide; and 30% to 95%, more preferably 40% to 90% of the delivery system.

By "tablet formulation" is meant the tablet made from the composition described herein, as well as the composition formulated in accordance with this disclosure but not in tablet form. The preferred tablet formulation of the present invention is in the form of a tablet.

Contemplated pesticides include those selected from the following classes, including mixtures thereof: herbicides, fungicides, bactericides, insecticides, nematocides, acaricides, and growth regulants.

20 Preferred dibasic and tribasic organic carboxylic acids include citric, fumaric, phthalic, maleic, malic, oxalic, adipic, glutaric, 2-methyl glutaric, succinic, and tartaric, or mixtures of any of them.

The term "dispersants" includes sodium, potassium, ammonium and calcium salts of naphthalene sulfonic acid-formaldehyde condensates; lithium, sodium, 25 potassium, calcium, and ammonium salts of lignosulfonates; sodium; potassium and ammonium salts of polyacrylates and carboxylates; sodium salts of maleic anhydride-isobutylene copolymers; and water soluble nonionic polymers such as polyvinyl-pyrrolidone, polyethylene oxides and cellulose derivatives. Preferred dispersants include the sodium, potassium, ammonium and calcium salts of naphthalene sulfonic acid-formaldehyde condensates, with the ammonium salts 30 more preferred.

Water-insoluble, cross-linked polyvinylpolypyrrolidone disintegrant refers to any of the generic, but is not limited to, crospovidone disintegrating agents.

35 The term "anionic wetting agent" includes, but is not limited to, salts of alkylbenzene sulfonates, alkyl and dialkyl naphthalene sulfonates, alkyl and

alcohol sulfates, sulfoalkylamides, carboxylates, alpha-olefin sulfonates and dialkyl sulfosuccinates. The term "nonionic wetting agent" includes acetylenic diols, ethylene oxide-propylene oxide copolymers, alkylphenol ethoxylates, fatty acid ethoxylates, alcohol ethoxylates, sorbitan fatty acid ester ethoxylates and castor oil ethoxylates. The preferred wetting agents are sodium dialkyl sulfosuccinates of which sodium diisobutyl sulfosuccinate, sodium diamyl sulfosuccinate and sodium dicyclohexyl sulfosuccinate are more preferred.

The internal desiccants that "chemically bind" water are those that actually undergo chemical reactions with water to form a new compound. An example of this type of material is CaO which reacts with water to form  $\text{Ca(OH)}_2$ . Other materials representative of those which react in this manner are magnesium oxide and boric anhydride.

The internal desiccants that "physically adsorb" water are those selected from the group consisting of highly-dispersed silicic acids such as silica gel; aluminum oxide; clays such as montmorillonite; and amorphous and crystalline aluminosilicates such as molecular sieves and zeolites. Combinations of these desiccants with those that form hydroxides and hydrates can be used. Kirk-Othmer's Encyclopedia of Chemical Technology (3rd ed., Vol. 8, p 115) describes desiccants suitable for use in the tablet formulation of this invention as Type 1 and Type 4 desiccants. Either type can be employed, singly or in combination, as long as the desiccant does not expand when it picks-up water. Such expansion causes the tablet to crack or crumble on long term storage.

Internal desiccants useful in the tablet formulation of this invention also include materials that chemically bind water, not in the sense of a chemical reaction that forms a hydroxide, but in the sense of a chemical reaction that produces a hydrate. Representative of useful desiccants that form hydrates are  $\text{CaSO}_4$ ,  $\text{NaOAc}$ ,  $\text{MgSO}_4$ ,  $\text{Na}_2\text{SO}_4$ ,  $\text{CaCl}_2$ , and  $\text{ZnSO}_4$ . Representative of the hydrate-forming reaction is that undergone by  $\text{CaCl}_2$  to form  $\text{CaCl}_2 \cdot \text{H}_2\text{O}$ . One or more desiccants from each group, the hydroxide-forming and the hydrate-forming, can be employed, alone or in combination, depending on the particular properties sought by the formulator. In any event, the desiccants employed in the tablets of this invention are not those of the water-adsorbing type employed in prior art tablet formulations. Kirk-Othmer's Encyclopedia of Chemical Technology (Third Edition, Vol. 8, page 115) further describes desiccants of the type contemplated for this invention as so-called Type 1 materials.

A preferred tablet formulation is one wherein component (b) is an ammonium, sodium or lithium carbonate or bicarbonate or mixture thereof, and the internal desiccant is selected from (A), (B) and a mixture of (A) and (B).

Also preferred is a tablet formulation wherein (b) is potassium carbonate or bicarbonate or mixture thereof, and the internal desiccant is (A).

Preferred pesticides are those having a melting point of at least about 100°C and solubility in pH 7 water at 20°C of no more than about 5% by weight.

Representatives of such pesticides are herbicides such as: acifluorfen, asulam, atrazine, bensulfuron, bentazon, bromacil, bromoxynil, chloramben, chlorimuron ethyl, chloroxuron, chlorsulfuron, chlortoluron, clomazone, cyanazine, dazomet, desmediphan, dicamba, dichlobenil, dichlorprop, diphenamid, dipropetryn, diuron, thiameturon, 2-[[[N-(4-methoxy-6-methyl-1,3,5-triazine-2-yl)-N-methylamino]carbonyl]amino]sulfonyl]benzoic acid, methyl ester, fenac, fenuron, fluometuron, fluridone, fomesafen, glyphosate, hexazinone, imazamethabenz, imazaquin, imazethapyr, ioxynil, isoproturon, isouron, isoxaben, karbutilate, lenacil, MCPA, MCPB, mefluidide, methabenzthiazuron, methazole, metribuzin, metsulfuron methyl, monuron, naptalam, neburon, nitralin, norflurazon, oryzalin, perfluidone, phenmedipham, picloram, prometryn, pronamide, propazine, pyrazon, siduron, simazine, sulfometuron methyl, tebuthiuron, terbacil, terbuthylazine, terbutryn, triclopyr, 2,4-D, 2,4-DB, triasulfuron, primisulfuron, 2-/3-(4,6-bis(difluoromethoxy)pyrimidin-2-yl)ureidosulfonyl]benzoic acid methyl ester, 5-pyrazolesulfonamide, N-[(4-methoxy-6-methyl-pyrimidine-2-yl)-amino-carbonyl]-4-methoxy-carbonyl-1-methyl-, N-[(4,6-dimethoxy-2-pyrimidinyl)-amino]carbonyl]-3-(ethylsulfonyl)-2-pyridinesulfonamide, 2-[[[(4,6-dimethoxy-2-pyrimidinyl)amino]carbonyl]amino]sulfonyl]-N,N-dimethyl-3-pyridine-carboxamide, methyl 2-[[[[4-ethoxy-6-(methylamino)-1,3,5-triazin-2-yl]-amino]carbonyl]amino]sulfonyl]benzoate, methyl 2-[[[(4,6-dimethoxy-2-pyrimidinyl)amino]carbonyl]amino]sulfonyl]-6-(trifluoromethyl)-3-pyridine-carboxylate, 2-(2-chloroethoxy)-N-[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)amino]carbonyl]benzenesulfonamide, methyl 2-[[[[4-(dimethyl-amino)-6-(2,2,2-trifluoroethoxy)-1,3,5-triazin-2-yl]amino]carbonyl]sulfonyl]-3-methyl-benzoate, sodium 2-chloro-6-[(4,6-dimethoxy-2-pyrimidinyl)thio]benzoate; fungicides such as: carbendazim, thiuram, dodine, chloroneb, cymoxanil, captan, folpet, thiophanatemethyl, thiabendazole, chlorothalonil, dichloran, captafol, iprodione, vinclozolin, kasugamycin, triadimenol, flutriafol, flusilazol,

hexaconazole or fenarimol; bactericides such as oxytetracycline dihydrate; acaricides such as: hexathiazox, oxythioquinox, dienochlor or cyhexatin; insecticides such as: carbofuran, carbaryl, thiodicarb or deltamethrin.

More preferred pesticides are hexazinone, 2,4-D, chlorsulfuron, 5  
sulfometuron methyl, chlorimuron ethyl, metsulfuron methyl, ethametsulfuron methyl, thifensulfuron methyl, tribenuron ethyl, bensulfuron methyl, primisulfuron, methyl 2-[[[(4,6-dimethoxy-2-pyrimidinyl)amino]carbonyl]-amino]sulfonyl]-6-(trifluoro-methyl)-3-pyridinecarboxylate, 2-(2-chloroethoxy)-*N*-[[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)amino]carbonyl]benzenesulfonamide, 10  
ethyl 5-[[[(4,6-dimethoxy-2-pyrimidinyl)amino]carbonyl]amino]sulfonyl]-1-methyl-1*H*-pyrazole-4-carboxylate, *N*-[[[(4,6-dimethoxy-2-pyrimidinylamino)carbonyl]-3-(ethylsulfonyl)-2-pyridinesulfonamide, 2-[[[(4,6-dimethoxy-2-pyrimidinyl)amino]carbonyl]amino]sulfonyl]-*N,N*-dimethyl-3-pyridine-carboxamide, methyl 2-[[[(4-(dimethylamino)-6-(2,2,2-trifluoroethoxy)-1,3,5- 15  
triazin-2-yl)amino]carbonyl]sulfonyl]-3-methylbenzoate, and sodium 2-chloro-6-[(4,6-dimethoxy-2-pyrimidinyl)thio]benzoate.

The most preferred pesticides are sulfonylurea herbicides such as chlorsulfuron, sulfometuron methyl, chlorimuron ethyl, metsulfuron methyl, ethametsulfuron methyl, thifensulfuron methyl, tribenuron ethyl, bensulfuron 20  
methyl, primisulfuron, methyl 2-[[[(4,6-dimethoxy-2-pyrimidinyl)amino]carbonyl]amino]sulfonyl]-6-(trifluoromethyl)-3-pyridinecarboxylate, 2-(2-chloroethoxy)-*N*-[[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)amino]carbonyl]benzenesulfonamide, ethyl 5-[[[(4,6-dimethoxy-2-pyrimidinyl)amino]carbonyl]amino]sulfonyl]-1-methyl-1*H*-pyrazole-4-carboxylate, *N*-[[[(4,6- 25  
dimethoxy-2-pyrimidinylamino)carbonyl]-3-(ethylsulfonyl)-2-pyridine-sulfonamide, 2-[[[(4,6-dimethoxy-2-pyrimidinyl)amino]carbonyl]amino]sulfonyl]-*N,N*-dimethyl-3-pyridinecarboxamide, and methyl 2-[[[(4-(dimethylamino)-6-(2,2,2-trifluoroethoxy)-1,3,5-triazin-2-yl)amino]carbonyl]-sulfonyl]-3-methylbenzoate.

### 30 DETAILS OF THE INVENTION

The most common method for applying water insoluble pesticides is as fine aqueous dispersions which are sprayed onto the field or crop using ground or aerial spray rigs. The tablets of this invention combine a high level of physical integrity with rapid break-up in water using minimal or no agitation while 35  
providing fine dispersions of active ingredient. Since the spray nozzles are

typically protected against clogging by 50 mesh screens (U.S. mesh size), the dispersions must be fine enough to pass through this size screen without plugging it. This ability is characteristic of pesticide dispersions delivered by the delivery system of this invention.

5 High physical integrity of the tablets is desirable so that the tablets themselves can withstand the tableting operation and survive handling, packaging and shipping without breaking. An axial breaking strength of greater than about  $9 \times 10^3$  newtons is generally necessary for a tablet to survive such treatment.

10 Rapid break-up in water is desirable for the convenience of the growers who require quick turnaround times for the preparation of the dispersions. Generally, the tablets of the invention disperse completely in less than 10 minutes, most in less than 5 minutes using even the cold water drawn from wells in the early spring.

15 It is substantially impossible to obtain rapid break-up of a tablet of substantially water-insoluble active ingredient in aqueous media without the use of effervescence. The reaction of the organic acid and carbonate or bicarbonate base affords carbon dioxide gas which aids in this respect.

A dispersant is required so that the particles of the active ingredient formed during the disintegration of the tablet remain separated in the cold, hard water.

20 The disintegrant allows the penetration of the water into the interior of the tablet through a wicking or swelling action. Common starch or cellulose-based disintegrants are unsuitable in agricultural applications as they typically form gels on the 50 mesh spray nozzle screens. Hence, a water insoluble cross-linked polyvinylpolypyrrolidone is used.

25 A wetting agent is required to control the size of the carbon dioxide bubbles formed during the reaction of the acid base. The wetting agent reduces the surface tension between the bubbles and the solid tablet resulting in the formation of smaller bubbles which readily detach from the tablet surface. As a consequence, the tablet remains submerged in the water for a longer period of time, thus improving contact of the entire tablet surface with water.

30 If a tablet floats immediately after being dropped in the water its top rapidly dries out and the reaction slows down there. This increases the time required for complete dispersion of active ingredient. When a tablet sinks, water wets the entire exterior of the tablet. Then, when the tablet floats to the surface (as a result of the buoyancy of the attached carbon dioxide bubbles when the tablet has

35



partially dispersed and become lighter) the top remains wet so that effervescent reaction continues. Dispersion times for active ingredients formulated as described herein are very much more rapid than in formulations that produce tablets designed for flotation. To ensure that the tablet will sink initially, inert ingredients are employed that produce a tablet with a density greater than that of water (specific gravity greater than 1.00).

Inert ingredients up to 99.9% of the total weight of the composition can be employed. Inert fillers such as sugar or clay can be added as long as they do not affect the chemical stability of the active ingredient(s). Materials such as glidants, anti-adherents, and lubricants can also be employed to facilitate production in the tablet press. The amounts and types of such ingredients will be readily determinable by one skilled in the tableting art given the disclosure herein.

The formulation ingredients are typically ground and mixed in a mill, e.g., an air or hammermill. The ground premix is passed through a 50 or 100 mesh (U.S.A. Standard Sieve Series) screen. The average particle size of the ground premix should be in the range of 5 to 15 microns. If it is much smaller, the tablet will be strong, but will not break up very fast. If the premix is much larger, the dispersion will not be fine enough to pass a wet screen test used to indicate whether the dispersion will clog the spray nozzle and protective screen discussed previously.

The tablets can be prepared using conventional tablet-making equipment. Their diameter can vary from about 1 cm or less, to 7.5 cm, depending on the tablet weight desired. Flat-faced, beveled-edge punches, with or without a breakline, produce attractive tablets.

To keep the tablet from sticking to the die or punch faces, a lubricant such as magnesium stearate or boric acid can be used. Such lubricants and anti-adherents can be brushed onto the die surface or incorporated into the formulation.

Tablets have been formed in a hydraulic press with a capacity of 18,000 kg of force. Pressures between about  $3.43 \times 10^7$  to  $6.86 \times 10^7$  pascals will produce strong tablets that break up rapidly. Break-up times are determined by dropping a tablet, typically 5 to 15 g into 1000 mL of water. The "end point" of final dispersion is easy to determine because the tablet floats to the surface as it loses weight shortly before it finally disperses.

The resultant dispersion is then poured through a nest of 50/100/200 mesh screens (300 mm, 150 mm, 75 mm holes, respectively). A qualitative judgment is then made about the amount of material that is retained on each screen. A good tablet will leave just a "trace" on the 200 mesh screen, and the larger screens will be free of residue.

The strength of the tablet can be measured by a tester such as the Erweka® Model TBH 28. The tablet is stood on end and the machine tip moves to the tablet along an axial path. The force to break the tablet in two is normally recorded in newtons (N). Good tablets normally have strengths in the range of  $8.896 \times 10^3$  to  $4.448 \times 10^4$  N.

#### EXAMPLE 1

The following ingredients were weighed out and milled for 1 min in a Tekmar® A-10 analytical laboratory mill. The premix was passed through a 50 mesh screen and blended well. A 15 g tablet, 4.34 cm in diameter, was made with a hand-operated hydraulic press at a pressure of  $525 \text{ kg/cm}^2$ .

Ingredient	Concentration, Weight %
Thifensulfuron methyl	55.3
Citric Acid	10.0
Sodium Bicarbonate	25.4
Lomar PWA® (ammonium salt of naphthalene sulfonic acid-formaldehyde condensate)	5.7
Polyplasdone XL-100® (polyvinylpyrrolidone)	1.22
Monawet MB-100® (sodium diisobutyl sulfosuccinate)	1.05
Magnesium Oxide	1.33

The fresh tablet took 2 min and 23 sec to dissolve in  $25^\circ\text{C}$  water with only a trace of solids on the screens. A second 15 g tablet was allowed to sit at room temperature for 3 days to allow the water in it to diffuse into the internal desiccant (MgO). The tablet was sealed in a tight jar and aged at  $45^\circ\text{C}$  for 3 weeks. This accelerated aging simulates about two years of storage at ambient conditions. After aging, the tablet took 3 min and 10 sec to dissolve in  $25^\circ\text{C}$  water. The wet screens had only a trace of solids on them.

EXAMPLE 2

The following ingredients were milled and tableted as in Example 1.

	<u>Ingredient</u>	<u>Concentration, Weight %</u>
5	Thifensulfuron methyl	52.7
	Citric Acid	9.5
	Sodium Bicarbonate	24.2
	Lomar PWA® (ammonium salt of naphthalene sulfonic acid-formaldehyde condensate)	5.45
10	Polyplasdone XL-100® (polyvinylpolypyrrolidone)	1.15
	Monawet MB-100® (sodium diisobutyl sulfosuccinate)	1.00
	Molecular Sieves	6.00

The fresh tablet took 2 min and 50 s to break-up in 25°C water. There was  
15 only a trace of solids on the screens. The second tablet was aged as in Example 1.  
After aging, the tablet took 2 min and 51 s to break-up. There was only a trace of  
solids on the wet screens.

EXAMPLES 3 TO 19

In the same manner employed for Example 1, tablets can be prepared using  
20 the active ingredients in the first column of Table 2 with one or more of the  
desiccants listed in the second and third columns, except that when the base  
employed is potassium carbonate or bicarbonate, the desiccant(s) employed  
therewith are to be selected solely from Column A.

TABLE 2


<u>Active Ingredient</u>	<u>Column (A)</u>	<u>Desiccant</u>
		<u>Column (B)</u>
3) hexazinone	selected from the	selected from the group
4) 2,4-D	group, CaO, MgO,	highly dispersed silicic
5) chlorsulfuron	B <sub>2</sub> O <sub>3</sub> , CaSO <sub>4</sub> ,	acids such as silica gel,
6) sulfometuron methyl	NaOAc, MgSO <sub>4</sub> ,	aluminum oxide, clays
7) chlorimuron ethyl	Na <sub>2</sub> SO <sub>4</sub> , CaCl <sub>2</sub> ,	such as
8) metsulfuron methyl	and ZnSO <sub>4</sub>	montmorillonite, and
9) ethametsulfuron methyl		amorphous aluminosilicates such as
10) tribenuron ethyl		molecular sieves and
11) bensulfuron methyl		zeolites
12) primisulfuron		
13) methyl 2-[[[(4,6-dimethoxy-2-pyrimidinyl)-amino]carbonyl]amino]sulfonyl]-6-(trifluoromethyl)-3-pyridinecarboxylate		
14) 2-(2-chloroethoxy)-N-[[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)amino]carbonyl]benzenesulfonamide		
15) ethyl 5-[[[(4,6-dimethoxy-2-pyrimidinyl)-amino]carbonyl]amino]sulfonyl]-1-methyl-1H-pyrazole-4-carboxylate		
16) N-[[[(4,6-dimethoxy-2-pyrimidinylamino)carbonyl]-3-(ethylsulfonyl)-2-pyridine-sulfonamide		
17) 2-[[[(4,6-dimethoxy-2-pyrimidinyl)amino]carbonyl]amino]sulfonyl]-N,N-dimethyl-3-pyridinecarboxamide		
18) methyl 2-[[[(4-(dimethylamino)-6-(2,2,2-trifluoroethoxy)-1,3,5-triazin-2-yl)amino]carbonyl]sulfonyl]-3-methylbenzoate		
19) sodium 2-chloro-6-[(4,6-dimethoxy-2-pyrimidinyl)thio]benzoate		

EXAMPLES 20 TO 37

By the general procedure of Example 1, tablet formulations can be made whereby the active ingredient pesticide is as described hereafter and the delivery system with its characteristic internal desiccant is as defined herein.


EXAMPLE 20

The pesticide, described in more detail in U.S. 4,127,405, is a compound of the formula:



wherein

$R_1$  is



$R_3$  and  $R_6$  are independently hydrogen, fluorine, chlorine, bromine, iodine, alkyl of 1-4 carbon atoms, alkoxy of 1-4 carbon atoms, nitro, trifluoromethyl, cyano,  $\text{CH}_3\text{S}(\text{O})_n-$  or  $\text{CH}_3\text{CH}_2\text{S}(\text{O})_n-$ ;

$R_4$  is hydrogen, fluorine, chlorine, bromine or methyl;

$R_5$  is hydrogen, fluorine, chlorine, bromine, methyl or methoxy;

$R_7$  is hydrogen, fluorine, chlorine, bromine, alkyl of 1-2 carbon atoms or alkoxy of 1-2 carbon atoms;

$R_8$  is hydrogen, methyl, chlorine or bromine;

$R_9$  and  $R_{10}$  are independently hydrogen, methyl, chlorine or bromine;

W and Q are independently oxygen or sulfur;

n is 0, 1 or 2;

X is hydrogen, chlorine, bromine, methyl, ethyl, alkoxy of 1-3 carbon atoms, trifluoromethyl,  $\text{CH}_3\text{S}-$  or  $\text{CH}_3\text{OCH}_2-$ ; and

5 Z is methyl or methoxy, or their agriculturally suitable salts.


#### EXAMPLE 21

The pesticide, described in more detail in U.S. 4,394,506, is a compound of the formula:

10 *N*-(heterocyclicaminocarbonyl)arylsulfonamides in which the aryl radical is substituted in the 2-position by a carboxy radical, ester, thioester, or amide thereof; e.g., *N*-[(4,6-dimethylpyrimidin-2-yl)aminocarbonyl]-methoxycarbonylbenzenesulfonamide or *N*-[(4,6-dimethoxy-1,3,5-triazin-2-yl)amino-carbonyl]-2-methoxycarbonylbenzenesulfonamide.

#### EXAMPLE 22

15 The pesticide, described in more detail in U.S. 4,481,029, is a compound of the formula:



20 wherein

W' is O or S;

A' is H, Cl, Br,  $\text{C}_1\text{-C}_4$  alkyl,  $\text{OCH}_3$ ,  $\text{NO}_2$  or  $\text{CF}_3$ ;

A is  $\begin{array}{c} \text{O} \\ \parallel \\ -\text{C}-\text{Q}-\text{R}^{\text{I}} \end{array}$  or  $\begin{array}{c} \text{T} \\ \parallel \\ -\text{C}-\text{R}^{\text{II}} \end{array}$  where

Q is O, S or  $\begin{array}{c} -\text{N}- \\ | \\ \text{R}_4 \end{array}$ ;


25

T is O or  $\begin{array}{c} \text{OR}^{\text{III}} \\ | \\ =\text{N} \end{array}$

where

$R^{III}$  is H,  $C_1$ - $C_4$  alkyl or  $C_3$ - $C_4$  alkenyl; when Q is O or S then

$R^I$  is  $C_1$ - $C_6$  alkyl  $C_3$ - $C_6$  alkenyl;  $C_3$ - $C_6$  alkynyl;  $C_2$ - $C_6$  alkyl substituted with 1-3 Cl, F or Br, or one of CN or  $OCH_3$ ;  $C_3$ - $C_6$  alkenyl substituted with 1-3 Cl;  $C_3$ - $C_6$  alkynyl substituted with Cl;  $C_5$ - $C_6$  cycloalkyl; cyclohexenyl; cyclohexyl substituted with 1-3  $CH_3$ ;  $C_4$ - $C_7$  cycloalkylalkyl or




10 where

$R_7$  and  $R_8$  are independently H, Cl,  $CH_3$  or  $OCH_3$ ;

$n$  is 0 or 1; and

$R_9$  is H or  $CH_3$ ;



15

where

20  $Z$  is N, CH or C-F;

$X$ =H, Cl,  $-CH_3$ ,  $-OCH_3$  or  $-OCH_2CH_3$ ;

$Y$ =H, Cl,  $C_1$ - $C_4$  substituted alkyl;

with the proviso that when  $X$  and  $Y$  are both H, then

$R^I$  and  $R^{II}$  are less than 5 carbons;

25  $X_1$ =H, Cl,  $OCH_3$ ,  $OCH_2CH_3$  or  $CH_3$ ;

$Y_1$ =H,  $OCH_3$  or  $CH_3$ ; and


$X_{III}=O$  or  $CH_2$  and further provided that when A contains greater than 5 carbon atoms, then Y contains  $\leq 4$  carbon atoms, and their agriculturally suitable salts;

all other substituents being as defined in U.S. 4,481,029.

5

### EXAMPLE 23

The pesticide, described in more detail in U.S. 4,435,205, is a compound of the formula:



10

where


W is O or S;

Q is O or  $NR_5$ ;

$R_1$  is  $C_1$ - $C_4$  alkyl,  $C_1$ - $C_4$  alkyl substituted with 1-3 atoms of F, Cl or Br,

15

$CH_2CH_2OCH_3$ ,  $CH_2CH_2CH_2OCH_3$  or



20

$R_2$  is H, F, Cl, Br,  $OCH_3$ ,  $NO_2$ ,  $CF_3$  or  $C_1$ - $C_2$  alkyl;

$R_3$  is H, F, Cl, Br or  $CH_3$ ;


$R_4$  is H,  $CH_3$  or  $OCH_3$ ;

$R_5$  is  $C_1$ - $C_4$  alkyl;

$R_6$  and  $R_7$  are independently H, F, Cl, Br,  $CH_3$ ,  $CF_3$ ,  $NO_2$  or  $OCH_3$ ;



A is



5

X is  $\text{NH}_2$ ,  $\text{N}(\text{CH}_3)_2$ ,  $\text{NHCH}_3$ ,  $\text{C}_1\text{-C}_4$  alkyl,  $\text{C}_1\text{-C}_4$  alkyl substituted with 1-3 atoms of F, Cl or Br,  $\text{CH}_2\text{OCH}_3$ ,  $\text{CH}_2\text{OCH}_2\text{CH}_3$ ,  $\text{C}_1\text{-C}_4$  alkoxy,  $\text{C}_1\text{-C}_2$  alkylthio,  $\text{C}_3\text{-C}_4$  alkenyloxy,  $\text{C}_3\text{-C}_4$  alkynyloxy,  $\text{OCH}_2\text{CH}_2\text{OCH}_3$  or  $\text{C}_2\text{-C}_4$  alkoxy substituted with 1-3 atoms of F, Cl or Br;

10

n is 1 or 2;

Y is H,  $\text{CH}_3$ ,  $\text{OCH}_3$  or Cl;

$\text{X}_1$  is O or  $\text{CH}_2$ ;

15

$\text{Y}_1$  is H,  $\text{CH}_3$ ,  $\text{OCH}_3$  or Cl;


$\text{X}_2$  and  $\text{Y}_2$  are independently  $\text{CH}_3$  or  $\text{OCH}_3$ ; and

Z is CH, N,  $\text{CCH}_3$ ,  $\text{CBr}$ ,  $\text{CCl}$ ,  $\text{CF}$ , Cl,  $\text{CC}_2\text{H}_5$ ,  $\text{CCH}_2\text{CH}_2\text{Cl}$  or  $\text{CCH}_2\text{CH}=\text{CH}_2$ .

#### EXAMPLE 24

20

The pesticide, described in more detail in U.S. 4,420,325, is a compound of the formula:



25

wherein

$R_1$  is F, Cl, Br,  $CF_3$ ,  $C_1$ - $C_3$  alkoxy,  $C_1$ - $C_3$  alkyl,  $NO_2$ ,  $CO_2R_4$ ,  $SO_2R_5$ ,  
 $SO_2NR_6R_7$ ,  $SO_2N(OCH_3)CH_3$ ,  $SO_2OCH_2CF_3$ ,  $OSO_2R_5$  or  $CH_2L$ ;

$L$  is  $SO_2NR_6R_7$ ,  $OCH_3$ ,  $OC_2H_5$ ,  $CO_2H_5$ ,  $CO_2CH_3$  or  $CO_2C_2H_5$ ;

$R_2$  is H, Cl, Br, F,  $CF_3$  or  $OCH_3$ ;

5  $R_4$  is  $C_1$ - $C_3$  alkyl,  $CH_2CH=CH_2$ ,  $CH_2CH_2Cl$  or  $CH_2CH_2OCH_3$ ;


$R_5$  is  $C_1$ - $C_3$  alkyl or  $CF_3$ ;

$R_6$  and  $R_7$  are independently  $C_1$ - $C_3$  alkyl;

$R_8$  is H or  $CH_3$ ;

$R_9$  is H or  $C_1$ - $C_3$  alkyl;

10  $R_3$  is



15

$W$  is O or S;

$X$  is  $CH_3$ ,  $OCH_3$  or Cl;

20  $Y$  is  $CH_3$ ,  $C_2H_5$ ,  $OCH_3$ ,  $OC_2H_5$ ,  $CH_2OCH_3$ ,  $NH_2$ ,  $NHCH_3$  or  $N(CH_3)_2$ ;

$Z$  is CH or N;

$X_1$  is H, Cl,  $CH_3$ ,  $OCH_3$  or  $OC_2H_5$ ;

$X_2$  is  $CH_3$ ,  $C_2H_5$ ,  $OCH_3$  or  $OC_2H_5$ ;

$X_3$  is  $CH_3$  or  $OCH_3$ ; and




25  $Y_1$  is  $CH_3$  or  $OCH_3$ ;

and their agriculturally suitable salts.


### EXAMPLE 25

The pesticide, described in more detail in U.S. 4,514,211, is a compound of the formula:

5




, or



10 wherein

Q is O, S, SO or SO<sub>2</sub>;

Q<sub>1</sub> is O, S or SO<sub>2</sub>;



15

R<sub>1</sub> is H or C<sub>1</sub>-C<sub>4</sub> alkyl;

$R_2$  is H or  $C_1$ - $C_4$  alkyl;

$R_3$  is H or  $CH_3$ ;

$R_4$  is H, Cl,  $CH_3$ ,  $CF_3$ ,  $OCH_3$ , Br, F,  $SCH_3$  or  $OCF_2H$ ;

$R_5$  is H,  $CH_3$ ,  $OCH_3$ , Cl, Br,  $NO_2$ ,  $CO_2R_7$ ,  $SO_2R_8$ ,  $OSO_2R_9$ ,

5  $SO_2NR_{10}R_{11}$ , F,  $CF_3$ ,  $SCH_3$ ,  $OCF_2H$  or  $SO_2N(OCH_3)CH_3$ ;

$R_6$  is H, Cl, Br or  $C_1$ - $C_4$  alkyl;

$R_6'$  is H,  $CH_3$ , Cl or Br;

$R_7$  is  $C_1$ - $C_3$  alkyl,  $CH_2CH=CH_2$ ,  $CH_2CH_2OCH_3$  or  $CH_2CH_2Cl$ ;

$R_8$  is  $C_1$ - $C_3$  alkyl;

10  $R_9$  is  $C_1$ - $C_3$  alkyl or  $CF_3$ ;






$R_{10}$  and  $R_{11}$  are independently  $C_1$ - $C_2$  alkyl;

$R_{12}$  is H or  $CH_3$ ;


W is O or S;

A is

15



or



20

X is H,  $CH_3$ ,  $OCH_3$ , Cl, F,  $OCF_2H$  or  $SCF_2H$ ;

Y is  $CH_3$ ,  $OCH_3$ ,  $OC_2H_5$ ,  $CH_2OCH_3$ ,  $NH_2$ ,  $NHCH_3$ ,  $N(CH_3)_2$ ,

$CH(OCH_3)_2$ ,  $CH(OCH_2CH_3)_2$ ,  $C_2H_5$ ,  $CF_3$ ,  $CH_2=CHCH_2O$ ,

25

$CH\equiv CCH_2O$ ,  $CF_3CH_2O$ ,  $OCH_2CH_2Cl$ ,  $OCH_2CH_2Br$ ,  $OCH_2CH_2F$ ,

CN,  $\text{CH}_2\text{OCH}_2\text{CH}_3$ ,  $\text{OCH}_2\text{CH}_2\text{OCH}_3$  or  $\text{GCF}_2\text{T}$  wherein G is O or S  
and T is H,  $\text{CHClF}$ ,  $\text{CHBrF}$ ,  $\text{CF}_2\text{H}$  or  $\text{CHFClF}_3$ ;

Z is CH, N,  $\text{CCH}_3$ ,  $\text{CC}_2\text{H}_5$ ,  $\text{CCl}$  or  $\text{CBr}$ ;

$\text{Y}_1$  is O or  $\text{CH}_2$ ;

5  $\text{X}_1$  is  $\text{CH}_3$ ,  $\text{OCH}_3$ ,  $\text{OC}_2\text{H}_5$  or  $\text{OCF}_2\text{H}$ ;


$\text{X}_2$  is  $\text{CH}_3$ ,  $\text{C}_2\text{H}_5$  or  $\text{CH}_2\text{CF}_3$ ;

$\text{Y}_2$  is  $\text{C}_2\text{H}_5$ ,  $\text{CH}_3$ ,  $\text{OCH}_3$ ,  $\text{OC}_2\text{H}_5$ ,  $\text{SCH}_3$  or  $\text{SC}_2\text{H}_5$ ; and

$\text{X}_3$  is  $\text{CH}_3$  or  $\text{OCH}_3$ .

#### EXAMPLE 26

10 The pesticide, described in more detail in U.S. 4,547,215, is a compound of the formula:




15 wherein

R is  $\text{C}_2\text{H}_5$  or  $\text{CH}(\text{CH}_3)_2$ ;

and their agriculturally suitable salts.

#### EXAMPLE 27

20 The pesticide, described in more detail in U.S. 4,548,638, is a compound of the formula:




wherein

25 R is  $\text{CO}_2\text{CH}_3$ ,  $\text{CO}_2\text{CH}_2\text{CH}_3$ ,  $\text{CO}_2\text{CH}_2\text{CH}_2\text{CH}_3$ ,  $\text{CO}_2\text{CH}_2\text{CH}=\text{CH}_2$ ,  
 $\text{CO}_2\text{CH}(\text{CH}_3)$ ,  $\text{CO}_2\text{CH}_2\text{CH}_2\text{Cl}$ ,  $\text{SO}_2\text{N}(\text{CH}_3)_2$  or  $\text{OSO}_2\text{CH}_3$ .

EXAMPLE 28

The pesticide, described in more detail in U.S. 4,479,821, is a compound of the formula:



wherein

A is a C<sub>1</sub>-C<sub>6</sub> alkyl radical which is substituted by C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> alkylthio, C<sub>1</sub>-C<sub>4</sub> alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub> alkylsulfonyl;

X is oxygen, sulfur, a sulfinyl or sulfonyl bridge;

Z is oxygen or sulfur;

m is 1 or 2;

R<sub>2</sub> is hydrogen, halogen, C<sub>1</sub>-C<sub>5</sub> alkyl, C<sub>2</sub>-C<sub>5</sub> alkenyl, C<sub>1</sub>-C<sub>4</sub> haloalkyl, or a radical -Y-R<sub>5</sub>, -COOR<sub>6</sub>, -NO<sub>2</sub> or -CO-NR<sub>7</sub>R<sub>8</sub>;

R<sub>3</sub> and R<sub>4</sub>, each independently of the other, are hydrogen, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> alkylthio, C<sub>1</sub>-C<sub>4</sub> haloalkyl, halogen or alkoxy-alkyl of at most 4 carbon atoms;


R<sub>5</sub> and R<sub>6</sub>, each independently of the other, are C<sub>1</sub>-C<sub>5</sub> alkyl, C<sub>2</sub>-C<sub>5</sub> alkenyl or C<sub>2</sub>-C<sub>6</sub> alkynyl;

R<sub>7</sub> and R<sub>8</sub>, each independently of the other, are hydrogen, C<sub>1</sub>-C<sub>5</sub> alkyl, C<sub>2</sub>-C<sub>5</sub> alkenyl or C<sub>2</sub>-C<sub>6</sub> alkynyl; and

Y is oxygen, sulfur, a sulfinyl or sulfonyl bridge, and salts of these compounds.


EXAMPLE 29

The pesticide, described in more detail in U.S. 4,566,898, is a compound of the formula:




EXAMPLE 30

The pesticide, described in more detail in U.S. 4,435,206, is a compound of the formula:



wherein

R is



R<sub>1</sub> is H, Cl, Br, F, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> alkylthio, NO<sub>2</sub>, CF<sub>3</sub>, COOR<sub>5</sub> or SO<sub>2</sub>NR<sub>6</sub>R<sub>7</sub>;

R<sub>2</sub> is H, Cl, Br or CH<sub>3</sub>;

R<sub>3</sub> and R<sub>4</sub> are independently H or CH<sub>3</sub>;

R<sub>5</sub> is C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>3</sub>-C<sub>6</sub> alkenyl, CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, CH<sub>2</sub>CH<sub>2</sub>OCH<sub>2</sub>CH<sub>3</sub>, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub> or CH<sub>2</sub>CH<sub>2</sub>Cl;

R<sub>6</sub> and R<sub>7</sub> are independently CH<sub>3</sub> or CH<sub>3</sub>CH<sub>2</sub>;

W is oxygen or sulfur;

X is CH<sub>3</sub>-OCH<sub>3</sub> or -OCH<sub>2</sub>CH<sub>3</sub>;

Y is H, Cl, CH<sub>3</sub>, CF<sub>3</sub>, -NHCH<sub>3</sub>, -N(CH<sub>3</sub>)<sub>2</sub>-, -CH<sub>2</sub>OR<sub>8</sub>, -CH<sub>2</sub>CH<sub>2</sub>OR<sub>8</sub>, -OCH<sub>2</sub>CF<sub>3</sub> or VR<sub>6</sub>;

Z is CH or N;

V is oxygen or sulfur;


R<sub>8</sub> is CH<sub>3</sub>, CH<sub>3</sub>CH<sub>2</sub>-, CH<sub>2</sub>CO<sub>2</sub>R<sub>8</sub>, -CH<sub>2</sub>CH<sub>2</sub>OR<sub>6</sub>, C(CH<sub>3</sub>)HCO<sub>2</sub>R<sub>8</sub> or CH<sub>2</sub>CH<sub>2</sub>CO<sub>2</sub>R<sub>8</sub>;

Y<sub>1</sub> is H, CH<sub>3</sub> or OCH<sub>3</sub>; and

$X_1$  is H, Cl,  $-\text{OCH}_3$ ,  $-\text{OCH}_2\text{CH}_3$  or  $\text{CH}_3$ ;  
and agricultural salts thereof.

### EXAMPLE 31

The pesticide, described in more detail in U.S. 4,514,212, is a compound of  
5 the formula:



and the salts thereof with amines, alkali metal or alkaline earth metal bases or  
10 with quaternary ammonium bases wherein:

Q is fluorine, fluoromethyl, chloromethyl, trichloromethyl, 1,2-dichloro-ethyl, 1,2-dibromoethyl, 1,2-dichloropropyl, 1,2-dibromopropyl, 1,2-dibromoisobutyl, 1,2-dichloro-1-methyl-ethyl or 1,2-dibromo-1-methylethyl;

15 X is oxygen, sulfur, a sulfinyl or sulfonyl bridge;

Z is oxygen or sulfur;

$R_2$  is hydrogen, halogen,  $\text{C}_1\text{-C}_5$  alkyl,  $\text{C}_2\text{-C}_5$  alkenyl,  $\text{C}_1\text{-C}_4$  haloalkyl, or a radical  $-\text{Y}-\text{R}_5$ ,  $-\text{COOR}_6$ ,  $-\text{NO}_2$  or  $-\text{CO}-\text{NR}_7-\text{R}_8$ ;

20  $R_3$  and  $R_4$ , each independently of the other, are hydrogen,  $\text{C}_1\text{-C}_4$  alkyl,  $\text{C}_1\text{-C}_4$  alkoxy,  $\text{C}_1\text{-C}_4$  alkylthio,  $\text{C}_1\text{-C}_4$  haloalkyl, halogen or alkoxyalkyl of at most 4 carbon atoms;

$R_5$  and  $R_6$ , each independently of the other, are  $\text{C}_1\text{-C}_5$  alkyl,  $\text{C}_2\text{-C}_5$  alkenyl or  $\text{C}_2\text{-C}_6$  alkynyl;

25  $R_7$  and  $R_8$ , each independently of the other, are hydrogen,  $\text{C}_1\text{-C}_5$  alkyl,  $\text{C}_2\text{-C}_5$  alkenyl or  $\text{C}_2\text{-C}_6$  alkynyl; and

Y is oxygen, sulfur, a sulfinyl or sulfonyl bridge.


### EXAMPLE 32

The pesticide, described in more detail in U.S. 4,478,635, is a compound of  
the formula:

30




23



wherein

X is a radical of the formula:

5



Y is C<sub>1</sub>-C<sub>3</sub> alkyl, C<sub>1</sub>-C<sub>3</sub> haloalkyl, C<sub>1</sub>-C<sub>3</sub> alkoxy, C<sub>1</sub>-C<sub>3</sub> haloalkoxy, C<sub>2</sub>-C<sub>3</sub> alkoxyalkyl, C<sub>1</sub>-C<sub>3</sub> alkylthio, halogen or -NR<sub>16</sub>R<sub>17</sub>;

10

Z is oxygen or sulfur;

R<sub>1</sub> is hydrogen, halogen, cyano, nitro, C<sub>1</sub>-C<sub>4</sub> haloalkyl, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, -CO-R<sub>6</sub>, -NR<sub>7</sub>R<sub>8</sub>, -S(O)<sub>m</sub>-C<sub>1</sub>-C<sub>4</sub> alkyl or -SO<sub>2</sub>R<sub>9</sub>;

R<sub>2</sub> is hydrogen, fluorine, chlorine, bromine, nitro, trifluoromethyl, -NR<sub>20</sub>R<sub>21</sub>, methyl, ethyl, methoxy, ethoxy or -S(O)<sub>m</sub>-C<sub>1</sub>-C<sub>4</sub> alkyl;

15

R<sub>3</sub> is hydrogen, fluorine, chlorine, bromine, amino, nitro or methoxy;

R<sub>6</sub> is hydrogen, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>3</sub> alkenyloxy, C<sub>3</sub>-C<sub>5</sub> alkynyloxy, C<sub>1</sub>-C<sub>4</sub> haloalkyl, C<sub>1</sub>-C<sub>5</sub> alkylthio, phenoxy, benzyloxy, -NR<sub>10</sub>R<sub>11</sub> or C<sub>1</sub>-C<sub>5</sub> alkoxy which is unsubstituted or substituted by 1 to 3 halogen atoms or C<sub>1</sub>-C<sub>3</sub> alkoxy;

20

R<sub>7</sub> is hydrogen, methoxy, ethoxy, C<sub>1</sub>-C<sub>4</sub> alkyl or -CO-R<sub>12</sub>;

R<sub>8</sub> is hydrogen or -CO-R<sub>12</sub>;

R<sub>9</sub> is an -O-R<sub>13</sub> or -NR<sub>14</sub>R<sub>15</sub> group;

R<sub>11</sub> is C<sub>1</sub>-C<sub>4</sub> alkyl which is unsubstituted or substituted by 1 to 3 halogen atoms, or is phenyl or benzyl;

25


R<sub>12</sub> is hydrogen, C<sub>1</sub>-C<sub>4</sub> alkyl or C<sub>1</sub>-C<sub>4</sub> alkoxy; and

m is 0, 1 or 2;

and  $R_4$  has the same meaning as  $R_2$ ;  $R_5$  has the same meaning as  $R_1$ ;  $R_{10}$ ,  $R_{11}$ ,  $R_{14}$  and  $R_{20}$  each have the same meaning as  $R_7$ ; and  $R_{12}$ ,  $R_{15}$ ,  $R_{16}$ ,  $R_{17}$  and  $R_{21}$  each have the same meaning as  $R_8$ .

**EXAMPLE 33**

5 The pesticide, described in more detail in U.S. 4,634,465, is a compound of the formula:



10 wherein

Z is oxygen or sulfur;

E is nitrogen or =C-;

$R_1$  is hydrogen, halogen, nitro,  $C_1$ - $C_4$  alkyl,  $C_1$ - $C_4$  haloalkyl,  $C_1$ - $C_4$  alkoxy,  $C_1$ - $C_4$  haloalkoxy,  $C_1$ - $C_4$  alkoxycarbonyl,  $C_1$ - $C_4$  alkylthio,  $C_1$ - $C_4$  alkylsulfinyl,  $C_1$ - $C_4$  alkylsulfonyl or  $C_2$ - $C_5$  alkoxyalkoxy;

$R_2$  is hydrogen,  $C_1$ - $C_4$  alkyl or  $C_1$ - $C_3$  alkoxy;

$R_3$  and  $R_4$ , each independently of the other, are hydrogen,  $C_1$ - $C_4$  alkyl,  $C_1$ - $C_4$  alkoxy,  $C_1$ - $C_4$  haloalkoxy,  $C_1$ - $C_4$  haloalkylthio,  $C_1$ - $C_4$  alkylthio, halogen,  $C_2$ - $C_5$  alkoxyalkyl,  $C_2$ - $C_5$  alkoxyalkoxy or  $-NR_5R_6$ , wherein  $R_5$  and  $R_6$  are hydrogen or  $C_1$ - $C_4$  alkyl; and

20 A is an unsubstituted or substituted bridge of 3 or 4 atoms which contains 1 or 2 oxygen, sulfur or nitrogen atoms and, together with the linking carbon atom, forms a non-aromatic 5- or 6-membered heterocyclic ring system, with the proviso that two oxygen atoms are separated by at least one carbon atom and that oxygen and sulfur atoms are only  
25 linked to each other if the sulfur atom takes the form of the -SO- or -SO<sub>2</sub>- group.


**EXAMPLE 34**

The pesticide, described in more detail in EPA-202,830, is:

30 2-[[N-(4-methoxy-6-methyl-1,3,5-triazin-2-yl)-N-methylamino-carbonyl]aminosulfonyl]benzoic acid, methyl ester.




EXAMPLE 35

The pesticide, described in more detail in EPA-237,292, is a compound of the formula:




wherein

J is



or



R is H or CH<sub>3</sub>;

R<sub>1</sub> is H or C<sub>1</sub>-C<sub>3</sub> alkyl;

R<sub>2</sub> is C<sub>1</sub>-C<sub>3</sub> alkyl or C<sub>1</sub>-C<sub>2</sub> alkoxy; or


R<sub>1</sub> and R<sub>2</sub> may be taken together to form -(CH<sub>2</sub>)<sub>n</sub>-, wherein n is 2, 3 or 4;

R<sub>3</sub> is H, Cl, F, Br, CH<sub>3</sub>, CF<sub>3</sub>, OCH<sub>3</sub> or COF<sub>2</sub>H; and

X is CH<sub>3</sub>, CH<sub>2</sub>F, CH<sub>2</sub>CH<sub>3</sub>, OCH<sub>3</sub>, OCH<sub>2</sub>CH<sub>3</sub>, Cl, OCF<sub>2</sub>H or CH<sub>2</sub>OCH<sub>3</sub>.

EXAMPLE 36

The pesticide, described in more detail in EPA-87,780, is a compound of the formula:




wherein

- 5       A represents a hydrogen atom, a C<sub>1</sub>-C<sub>8</sub> alkyl group or a phenyl group which may be substituted with C<sub>1</sub>-C<sub>8</sub> alkyl groups, halogen atoms or nitro groups; B and C represent independently hydrogen atoms, halogen atoms, nitro groups, C<sub>1</sub>-C<sub>8</sub> alkyl groups, arylalkyl groups, C<sub>1</sub>-C<sub>8</sub> alkoxy groups; haloalkyl groups, -CO<sub>2</sub>R [where R is a hydrogen atom, a C<sub>1</sub>-C<sub>8</sub> alkyl group, an allyl group or a propargyl group), -CONR<sub>1</sub>R<sub>2</sub> (where R<sub>1</sub> is a hydrogen atom, a C<sub>1</sub>-C<sub>8</sub> alkyl group or a phenyl group, R<sub>2</sub> is a hydrogen atom or a C<sub>1</sub>-C<sub>8</sub> alkyl group, or R<sub>1</sub> and R<sub>2</sub> taken together may represent -(CH<sub>2</sub>)<sub>m</sub>- (m is 4, 5 or 6), -CH<sub>2</sub>CH<sub>2</sub>OCH<sub>2</sub>CH<sub>2</sub>-, or -CH<sub>2</sub>CH<sub>2</sub>N(CH<sub>3</sub>)CH<sub>2</sub>CH<sub>2</sub>-],
- 10       -S(O)<sub>n</sub>R<sub>3</sub> (where R<sub>3</sub> is a C<sub>1</sub>-C<sub>8</sub> alkyl group, a phenyl group or an arylalkyl group and n is 0, 1 or 2), -SO<sub>2</sub>NR<sub>4</sub>R<sub>5</sub> [where R<sub>4</sub> is a C<sub>1</sub>-C<sub>8</sub> alkyl group, R<sub>5</sub> is a hydrogen atom or a C<sub>1</sub>-C<sub>8</sub> alkyl group, or R<sub>4</sub> and R<sub>5</sub> taken together may represent -(CH<sub>2</sub>)<sub>p</sub>- (p is 4, 5 or 6), -CH<sub>2</sub>CH<sub>2</sub>OCH<sub>2</sub>CH<sub>2</sub>- or -CH<sub>2</sub>CH<sub>2</sub>N(CH<sub>3</sub>)CH<sub>2</sub>CH<sub>2</sub>-] or a phenyl group which may be substituted with C<sub>1</sub>-C<sub>8</sub> alkyl groups, halogen atoms or nitro groups, D represents a hydrogen atom or a C<sub>1</sub>-C<sub>8</sub> alkyl group; X and Y represent independently hydrogen atoms, halogen atoms, C<sub>1</sub>-C<sub>8</sub> alkyl groups, C<sub>1</sub>-C<sub>8</sub> alkoxy groups, C<sub>1</sub>-C<sub>8</sub> alkoxyalkyl groups, -CF<sub>3</sub> groups, C<sub>1</sub>-C<sub>8</sub> haloalkoxy groups, alkylamino
- 15       R  
/
- 20       groups, dialkylamino groups, -OCHCO<sub>2</sub>R<sub>7</sub> (where R<sub>6</sub> and R<sub>7</sub> each represent hydrogen atoms or C<sub>1</sub>-C<sub>8</sub> alkyl groups) or either X or Y may form a five-membered ring containing an oxygen atom together with X; and X represents a nitrogen atom or C-R<sub>8</sub> (where R<sub>8</sub> represents a hydrogen atom, a haloalkyl group or may
- 25
- 30

form a five-membered ring containing an oxygen atom together with X or Y).

### EXAMPLE 37

The pesticide, described in more detail in U.S. 4,710,221, is a compound of the formula:



wherein

- 10 R is H or CH<sub>3</sub>;  
 R<sub>1</sub> is C<sub>1</sub>-C<sub>3</sub> alkyl, C<sub>3</sub>-C<sub>4</sub> alkoxyalkyl, C<sub>2</sub>-C<sub>4</sub> haloalkyl, C<sub>3</sub>-C<sub>4</sub> alkenyl or C<sub>3</sub>-C<sub>4</sub> alkynyl;  
 R<sub>2</sub> is C<sub>2</sub>-C<sub>6</sub> alkoxy, C<sub>3</sub>-C<sub>6</sub> cycloalkoxy, C<sub>4</sub>-C<sub>6</sub> cycloalkylalkoxy, C<sub>1</sub>-C<sub>6</sub> haloalkoxy, C<sub>2</sub>-C<sub>6</sub> alkenyloxy, C<sub>2</sub>-C<sub>6</sub> haloalkenyloxy, C<sub>3</sub>-C<sub>6</sub> alkynyloxy, C<sub>3</sub>-C<sub>6</sub> haloalkynyloxy, C<sub>2</sub>-C<sub>4</sub> alkoxyalkoxy, C<sub>2</sub>-C<sub>4</sub> haloalkoxyalkoxy, C<sub>2</sub>-C<sub>4</sub> alkylthioalkoxy, C<sub>2</sub>-C<sub>4</sub> haloalkylthio-  
 15 alkoxy, C<sub>2</sub>-C<sub>4</sub> alkylsulfinylalkoxy, C<sub>2</sub>-C<sub>4</sub> haloalkylsulfinylalkoxy, C<sub>2</sub>-C<sub>4</sub> alkylsulfonylalkoxy, C<sub>2</sub>-C<sub>4</sub> haloalkylsulfonylalkoxy, C<sub>2</sub>-C<sub>4</sub> cyanoalkoxy, OCH<sub>2</sub>C(O)CH<sub>3</sub>, OCH<sub>2</sub>CH<sub>2</sub>C(O)CH<sub>3</sub>, C<sub>2</sub>-C<sub>4</sub> aminoalkoxy, C<sub>1</sub>-C<sub>8</sub> alkylthio, C<sub>3</sub>-C<sub>6</sub> cycloalkylthio, C<sub>4</sub>-C<sub>6</sub> cycloalkylalkylthio, C<sub>1</sub>-C<sub>8</sub> haloalkylthio, C<sub>2</sub>-C<sub>6</sub> alkenylthio, C<sub>2</sub>-C<sub>6</sub> haloalkenylthio, C<sub>3</sub>-C<sub>6</sub> alkynylthio, C<sub>3</sub>-C<sub>6</sub> haloalkynylthio, C<sub>2</sub>-C<sub>4</sub> alkoxyalkylthio, C<sub>2</sub>-C<sub>4</sub> haloalkoxyalkylthio, C<sub>2</sub>-C<sub>4</sub> alkylthio-  
 20 alkylthio, C<sub>2</sub>-C<sub>4</sub> haloalkylthioalkylthio, C<sub>2</sub>-C<sub>4</sub> cyanoalkylthio, SCH<sub>2</sub>C(O)CH<sub>3</sub>, SCH<sub>2</sub>CH<sub>2</sub>C(O)CH<sub>3</sub>, C<sub>2</sub>-C<sub>4</sub> aminoalkylthio, SC<sub>6</sub>H<sub>5</sub>, SCH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>, C<sub>1</sub>-C<sub>8</sub> alkylsulfinyl, C<sub>3</sub>-C<sub>6</sub> cycloalkylsulfinyl, C<sub>4</sub>-C<sub>6</sub> cycloalkylalkylsulfinyl, C<sub>1</sub>-C<sub>8</sub> haloalkylsulfinyl, C<sub>2</sub>-C<sub>6</sub> alkenylsulfinyl, C<sub>2</sub>-C<sub>6</sub> haloalkenylsulfinyl, C<sub>3</sub>-C<sub>6</sub> alkynylsulfinyl, C<sub>3</sub>-C<sub>6</sub> haloalkynylsulfinyl, C<sub>2</sub>-C<sub>4</sub> alkoxyalkylsulfinyl, C<sub>2</sub>-C<sub>4</sub> haloalkoxyalkylsulfinyl, C<sub>2</sub>-C<sub>4</sub> cyanoalkylsulfinyl, S(O)CH<sub>2</sub>C(O)CH<sub>3</sub>,  
 25 S(O)CH<sub>2</sub>CH<sub>2</sub>C(O)CH<sub>3</sub>, C<sub>2</sub>-C<sub>4</sub> aminoalkylsulfinyl, C<sub>2</sub>-C<sub>8</sub> alkylsulfonyl, C<sub>3</sub>-C<sub>6</sub> cycloalkylsulfonyl, C<sub>4</sub>-C<sub>6</sub> cycloalkylalkyl-
- 30

sulfonyl, C<sub>1</sub>-C<sub>8</sub> haloalkylsulfonyl, C<sub>2</sub>-C<sub>6</sub> alkenylsulfonyl, C<sub>2</sub>-C<sub>6</sub>  
 haloalkenylsulfonyl, C<sub>3</sub>-C<sub>6</sub> alkynylsulfonyl, C<sub>3</sub>-C<sub>6</sub> haloalkynyl-  
 sulfonyl, C<sub>2</sub>-C<sub>4</sub> alkoxyalkylsulfonyl, C<sub>2</sub>-C<sub>4</sub> haloalkoxyalkylsulfonyl,  
 C<sub>2</sub>-C<sub>4</sub> cyanoalkylsulfonyl, SO<sub>2</sub>CH<sub>2</sub>C(O)CH<sub>3</sub>,  
 5 SO<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>C(O)CH<sub>3</sub>, C<sub>2</sub>-C<sub>4</sub> aminoalkylsulfonyl, CH<sub>2</sub>F, CH<sub>2</sub>Cl,  
 CHCl<sub>2</sub>, CH<sub>2</sub>Br, CHBr<sub>2</sub>, C<sub>2</sub>-C<sub>6</sub> alkyl substituted with 1-3 atoms of F,  
 Cl or Br, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> haloalkenyl, C≡CH, C<sub>2</sub>-C<sub>6</sub>  
 haloalkynyl, OC(O)C<sub>1</sub>-C<sub>4</sub> alkyl, CH<sub>2</sub>C(O)NR<sub>a</sub>R<sub>b</sub>, NHCH<sub>3</sub>, NR<sub>b</sub>R<sub>c</sub>  
 or C<sub>1</sub>-C<sub>4</sub> alkyl substituted with C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>3</sub>-C<sub>4</sub> cycloalkoxy,  
 10 cyclopropylmethoxy, C<sub>1</sub>-C<sub>4</sub> haloalkoxy, C<sub>2</sub>-C<sub>4</sub> alkenyloxy, C<sub>2</sub>-C<sub>4</sub>  
 haloalkenyloxy, C<sub>3</sub>-C<sub>4</sub> alkynyloxy, C<sub>3</sub>-C<sub>4</sub> haloalkynyloxy, C<sub>2</sub>-C<sub>4</sub>  
 alkoxyalkoxy, C<sub>2</sub>-C<sub>4</sub> aminoalkoxy, C<sub>1</sub>-C<sub>4</sub> alkylcarbonyloxy, C<sub>1</sub>-C<sub>4</sub>  
 haloalkylcarbonyloxy, C<sub>1</sub>-C<sub>4</sub> carbamoyloxy, C<sub>1</sub>-C<sub>4</sub> alkoxy-  
 carbonyloxy, OH, OP(O)(OC<sub>1</sub>-C<sub>2</sub> alkyl)<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub> alkylsulfonyloxy,  
 15 C<sub>1</sub>-C<sub>2</sub> haloalkylsulfonyloxy, OSi(CH<sub>3</sub>)<sub>3</sub>, OSi(CH<sub>3</sub>)<sub>2</sub>C(CH<sub>3</sub>)<sub>3</sub>, C<sub>1</sub>-C<sub>4</sub>  
 alkylthio, C<sub>3</sub>-C<sub>4</sub> cycloalkylthio, cyclopropylmethylthio, C<sub>1</sub>-C<sub>4</sub>  
 haloalkylthio, C<sub>2</sub>-C<sub>4</sub> alkenylthio, C<sub>2</sub>-C<sub>4</sub> haloalkenylthio, C<sub>3</sub>-C<sub>4</sub>  
 alkynylthio, C<sub>3</sub>-C<sub>4</sub> haloalkynylthio, C<sub>2</sub>-C<sub>4</sub> alkoxyalkylthio, C<sub>2</sub>-C<sub>4</sub>  
 aminoalkylthio, SH, SP(O)(OC<sub>1</sub>-C<sub>2</sub> alkyl)<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub> alkylsulfinyl,  
 20 C<sub>3</sub>-C<sub>4</sub> cycloalkylsulfonyl, cyclopropylmethylsulfinyl, C<sub>1</sub>-C<sub>4</sub>  
 haloalkylsulfinyl, C<sub>2</sub>-C<sub>4</sub> alkenylsulfinyl, C<sub>2</sub>-C<sub>4</sub> haloalkenylsulfinyl,  
 C<sub>3</sub>-C<sub>4</sub> alkynylsulfinyl, C<sub>3</sub>-C<sub>4</sub> haloalkynylsulfinyl, C<sub>2</sub>-C<sub>4</sub>  
 alkoxyalkylsulfinyl, C<sub>2</sub>-C<sub>4</sub> aminoalkylsulfinyl, C<sub>1</sub>-C<sub>4</sub> alkylsulfonyl,  
 C<sub>3</sub>-C<sub>4</sub> cycloalkylsulfonyl, cyclopropylmethylsulfonyl, C<sub>1</sub>-C<sub>4</sub>  
 25 haloalkylsulfonyl, C<sub>2</sub>-C<sub>4</sub> alkenylsulfonyl, C<sub>2</sub>-C<sub>4</sub> haloalkenylsulfonyl,  
 C<sub>3</sub>-C<sub>4</sub> alkynylsulfonyl, C<sub>3</sub>-C<sub>4</sub> haloalkynylsulfonyl, C<sub>2</sub>-C<sub>4</sub>  
 alkoxyalkylsulfonyl or C<sub>2</sub>-C<sub>4</sub> aminoalkylsulfonyl;

R<sub>a</sub> and R<sub>b</sub> are independently H or C<sub>1</sub>-C<sub>3</sub> alkyl;

30 R<sub>c</sub> is C<sub>2</sub>-C<sub>4</sub> alkyl, cyclopropylmethyl, C<sub>2</sub>-C<sub>4</sub> cyanoalkyl, CH<sub>2</sub>C(O)CH<sub>3</sub>,  
 CH<sub>2</sub>CH<sub>2</sub>C(O)CH<sub>3</sub>, C<sub>10</sub>-C<sub>4</sub> haloalkyl, C<sub>3</sub>-C<sub>4</sub> alkenyl, C<sub>3</sub>-C<sub>4</sub>  
 haloalkenyl, C<sub>3</sub>-C<sub>4</sub> alkynyl, C<sub>3</sub>-C<sub>4</sub> haloalkynyl, C<sub>1</sub>-C<sub>4</sub> alkyl  
 substituted with C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> alkylthio, C<sub>1</sub>-C<sub>4</sub> alkylsulfinyl,  
 C<sub>1</sub>-C<sub>4</sub> alkylsulfonyl, OH, NH<sub>2</sub>, NHCH<sub>3</sub> or N(CH<sub>3</sub>)<sub>2</sub>;

X is CH<sub>3</sub>, OCH<sub>3</sub>, OC<sub>2</sub>H<sub>5</sub>, Cl or Br;

Y is C<sub>1</sub>-C<sub>2</sub> alkyl, C<sub>1</sub>-C<sub>2</sub> alkoxy, OCH<sub>2</sub>CH<sub>2</sub>F, OCH<sub>2</sub>CHF<sub>2</sub>, OCH<sub>2</sub>CF<sub>3</sub>,  
NHCH<sub>3</sub> or N(CH<sub>3</sub>)<sub>2</sub>; and

Z is CH or N; and

their agriculturally suitable salts.

CLAIMS

What is claimed is:

- 5 1. A tablet formulation consisting essentially of by total weight of the formulated composition:
  - (i) about 0.1% to 75% of a pesticide;
  - (ii) about 25% to 99.9% of a delivery system characterized by a panel of components complementary to the pesticide of (i) having the following
 10 components:
    - (a) about 5% to 75% of a dibasic or tribasic organic carboxylic acid or a mixture thereof;
    - (b) about 5% to 75% of an ammonium, lithium, sodium or potassium carbonate or bicarbonate or a mixture thereof;
    - 15 (c) about 0.5% to 20% of a dispersant;
    - (d) about 0.1% to 5% of water-insoluble cross-linked polyvinylpyrrolidone;
    - (e) about 0.1% to 5% of an anionic or nonionic wetting agent; and
    - (f) about 1% to 20% of an internal desiccant being selected from the
 20 group:
      - (A) one or a mixture of desiccants that chemically bind water, and
      - (B) one or a mixture of desiccants that physically adsorb water;

the desiccant being (A) when (b) is potassium carbonate or potassium bicarbonate.
- 25 2. A tablet formulation according to Claim 1 wherein (b) is an ammonium, sodium or lithium carbonate or bicarbonate or mixture thereof, and the internal desiccant is selected from (A), (B) and a mixture of (A) and (B).
- 3 3. A tablet formulation according to Claim 1 wherein (b) is potassium carbonate or bicarbonate or mixture thereof, and the internal desiccant is (A).
- 4 4. A tablet formulation according to Claim 1 in the form of a tablet.
- 30 5. A tablet formulation according to Claim 2 in the form of a tablet.
- 6 6. A tablet formulation according to Claim 3 in the form of a tablet.
- 7 7. A tablet formulation according to any one of Claims 1 to 6 wherein the pesticide is a sulfonylurea herbicide selected from the group consisting of
 

chlorsulfuron, sulfometuron methyl, chlorimuron ethyl, metsulfuron methyl,
 35 ethametsulfuron methyl, thifensulfuron methyl, tribenuron ethyl, bensulfuron methyl, primisulfuron, methyl 2-[[[(4,6-dimethoxy-2-pyrimidinyl)amino]-



carbonyl]amino)sulfonyl]-6-(trifluoro-methyl)-3-pyridinecarboxylate, 2-(2-chloroethoxy)-*N*-[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)amino]carbonyl]-benzenesulfonamide, ethyl 5-[[(4,6-dimethoxy-2-pyrimidinyl)amino]carbonyl]-amino)sulfonyl]-1-methyl-1*H*-pyrazole-4-carboxylate, *N*-[[(4,6-dimethoxy-2-pyrimidinylamino)carbonyl]-3-(ethylsulfonyl)-2-pyridinesulfonamide, 2-[[(4,6-dimethoxy-2-pyrimidinyl)amino]carbonyl]amino)sulfonyl]-*N,N*-dimethyl-3-pyridinecarboxamide, and methyl 2-[[(4-(dimethylamino)-6-(2,2,2-trifluoroethoxy)-1,3,5-triazin-2-yl)amino]carbonyl]sulfonyl]-3-methylbenzoate.

8. A tablet formulation according to Claim 7 wherein the pesticide is a  
10 sulfonylurea herbicide selected from the group consisting of thifensulfuron methyl, tribenuron ethyl, and bensulfuron methyl.

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 92/11305

<b>I. CLASSIFICATION OF SUBJECT MATTER</b> (If several classification symbols apply, indicate all) <sup>6</sup>		
According to International Patent Classification (IPC) or to both National Classification and IPC Int.Cl. 5 A01N25/34;                      A01N47/36;                      A01N43/64;                      A01N39/04		
<b>II. FIELDS SEARCHED</b>		
Minimum Documentation Searched <sup>7</sup>		
Classification System	Classification Symbols	
Int.Cl. 5	A01N	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched <sup>8</sup>		
<b>III. DOCUMENTS CONSIDERED TO BE RELEVANT<sup>9</sup></b>		
Category <sup>10</sup>	Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup>	Relevant to Claim No. <sup>13</sup>
X	EP,A,0 391 851 (CIBA-GEIGY) 10 October 1990 see column 1, line 1 - line 24 see column 2, line 1 - line 20 see column 2, line 36 - column 4, line 16 see column 5, line 2 - column 9, line 55	1,2,4,5, 7,8
Y	& CA,A,2 013 918 cited in the application ---	3,6
Y	PATENT ABSTRACTS OF JAPAN vol. 9, no. 1 (C-259)(1724) 5 January 1985 & JP,A,59 155 311 (NISSAN-) 4 September 1984 see abstract ---	3,6
-/-		
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p><sup>10</sup> Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&amp;" document member of the same patent family</p> </div> </div>		
<b>IV. CERTIFICATION</b>		
Date of the Actual Completion of the International Search <div style="text-align: center;">18 MARCH 1993</div>	Date of Mailing of this International Search Report <div style="text-align: center;">20. 04. 93</div>	
International Searching Authority <div style="text-align: center;">EUROPEAN PATENT OFFICE</div>	Signature of Authorized Officer <div style="text-align: center;">LAMERS W.</div>	

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		Relevant to Claim No.
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	
Y	WO,A,9 000 007 (E.I.DU PONT DE NEMOURS) 11 January 1990 see the whole document ----	1,2,4,5, 7,8
Y	GB,A,2 242 130 (INFOWISE) 25 September 1991 see page 4, line 2 - line 19 see page 6; example B see page 7, line 7 - line 12 ----	1,2,4,5, 7,8
P,X	EP,A,0 488 660 (SUMITOMO CHEMICAL) 3 June 1992 see page 2, line 10 - page 7, line 5 see page 7, line 21 - line 24 ----	1-8
A	EP,A,0 415 688 (AECI) 6 March 1991 see the whole document ----	1-8
A	EP,A,0 367 887 (CIBA-GEIGY) 16 May 1990 see the whole document ----	1-8
A	US,A,4 180 467 (J.B.BARTH) 25 December 1979 see column 1, line 11 - line 26 see column 1, line 65 - column 2, line 22 ----	1-8
A	DE,A,2 710 107 (HOYER) 14 September 1978 see page 2; claims 9,10 see page 8, paragraph 3 - page 9, paragraph 3 -----	1-8

**ANNEX TO THE INTERNATIONAL SEARCH REPORT  
ON INTERNATIONAL PATENT APPLICATION NO.**

US 9211305  
SA 68875

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report.  
The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

18/03/93

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0391851	10-10-90	AU-B- 625467	09-07-92
		AU-A- 5304290	11-10-90
		CA-A- 2013918	07-10-90
		JP-A- 2290801	30-11-90
WO-A-9000007	11-01-90	AU-B- 626551	06-08-92
		AU-A- 3691189	23-01-90
		CN-A- 1039948	28-02-90
		EP-A- 0422026	17-04-91
		JP-T- 3505451	28-11-91
		US-A- 5180587	19-01-93
GB-A-2242130	25-09-91	None	
EP-A-0488660	03-06-92	JP-A- 5000903	08-01-93
EP-A-0415688	06-03-91	AU-A- 6204390	07-03-91
		CA-A- 2024291	01-03-91
		JP-A- 3163004	15-07-91
EP-A-0367887	16-05-90	US-A- 4933000	12-06-90
		AU-B- 625412	09-07-92
		AU-A- 2588688	08-02-90
		JP-A- 2048506	19-02-90
US-A-4180467	25-12-79	None	
DE-A-2710107	14-09-78	None	